


**MICROBIOLOGICAL IMAGE**

# The microPIXE technique to understand the distribution of heavy metals in arbuscular mycorrhizal symbiosis

## Uso de la técnica de microPIXE para comprender la distribución de metales pesados en la simbiosis micorrízica arbuscular

M. Statello<sup>a,b</sup>, R.P. Colombo<sup>a,b,\*</sup>, E.M. de la Fournière<sup>c</sup>, M.E. Debray<sup>c</sup>, A.M. Godeas<sup>a,b</sup>, V.A. Silvani<sup>a,b</sup>

<sup>a</sup> Instituto de Biodiversidad y Biología Experimental y Aplicada, UBA-CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina

<sup>b</sup> Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

<sup>c</sup> Gerencia Investigación y Aplicaciones, Comisión Nacional de Energía Atómica, CAC, San Martín, Buenos Aires, Argentina

Received 8 September 2023; accepted 31 December 2023


Available online 14 May 2024

Soil colloids tend to retain heavy metals (HMs), potentially reaching toxic concentrations. Phytoremediation, a decontamination technique, involves plants and their associated microorganisms to mitigate the toxicity of HMs in the environment. Most plant species employed in phytoremediation establish symbiotic associations with arbuscular mycorrhizal fungi (AMF). AMF can adsorb or uptake HMs through their hyphal network, translocating them towards the host plant<sup>1</sup>. Particle-induced X-ray emission with a micro-focused beam (microPIXE) enables the micrometric mapping of elements in a sample<sup>2</sup>, helping to determine the concentration and distribution of HMs. In this work, we used microPIXE to analyze the distribution and accumulation of copper (Cu) in structures of the AMF *Rhizophagus intraradices* GC3 strain associated to Ri T-DNA carrot roots under *in vitro* condi-

tion (*Banco de Glomeromycota in vitro*)<sup>3</sup>. This would enable us to understand the role of AMF in HM-polluted soils. An *in vitro* assay was performed using bi-compartmentalized Petri plates. The root compartment (RC) contained solid minimal medium (MM) with transformed carrot roots colonized by GC3. The hyphal compartment (HC), where only AMF developed, contained sterile soil artificially contaminated with Cu (300 mg/kg)<sup>4</sup>. After three months, roots and intraradical vesicles from RC and extraradical mycelium from HC were sampled for microPIXE analysis. The map of Ca, a major element in biological systems, helped us to visualize the biological structures and identify other elements; P and Mn (a MM compound) were selected as AMF structure indicators. P, Mn and Cu, were mainly detected in vesicles, and to a lesser extent in colonized roots (considering the dimensions of the structures), while P and Mn were mostly localized in the extraradical mycelium (Fig. 1). These results showed significant Cu accumulation in intraradical fungal structures but not in root cells, suggesting a potential mech-

\* Corresponding author.

E-mail address: [colomboroxanap@gmail.com](mailto:colomboroxanap@gmail.com) (R.P. Colombo).



**Figure 1** 2D X-ray emission distribution maps of Ca, P, Mn and Cu from the extraradical mycelium and vesicles of the arbuscular mycorrhizal fungus *Rhizophagus intraradices* GC3 strain and a cross section of a mycorrhized carrot root. MicroPIXE conditions: 50 MeV  $^{16}\text{O}^{5+}$  beam scan size  $250\ \mu\text{m} \times 250\ \mu\text{m}$  and  $200\ \mu\text{m} \times 200\ \mu\text{m}$ , respectively. Spot size  $3\ \mu\text{m} \times 3\ \mu\text{m}$ . The colour scale, from blue to red, represents the increase in the concentration of the chemical elements.

anism of toxicity alleviation in the host plant, and laying the foundation for future studies focusing on Mn.

## Funding

This work was supported by Universidad de Buenos Aires (UBA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Agencia Nacional de Promoción Científica y Tecnológica (ANPyCT) and Comisión Nacional de Energía Atómica (CNEA).

## Conflict of interest

Authors have no conflict of interest.

## References

1. Cabral L, Soares CRFS, Giachini AJ, Siqueira JO. Arbuscular mycorrhizal fungi in phytoremediation of contaminated areas by trace elements: mechanisms and major benefits of their applications. *World J Microbiol Biotechnol*. 2015;31:1655–64, <http://dx.doi.org/10.1007/s11274-015-1918-y>.
2. de la Fournière EM, Vega NA, Muller NA, Debray ME. MicroPIXE analysis of removal of aqueous U(VI) by *S. intermedia* and *P. striatipes* in the presence of Th (IV). *Nucl Inst Meth Phys Res B*. 2020;477:87–90, <http://dx.doi.org/10.1016/j.nimb.2019.11.001>.
3. <https://bgiv.com.ar/> [last accessed: 13.11.23].
4. Silvani VA, Statello M, Scorza MV, Pèrgola M, Colombo RP, Godeas AM. A novel *in vitro* methodology to cultivate arbuscular mycorrhizal fungi combining soil and synthetic media. *Symbiosis J*. 2019;79:163–70, <http://dx.doi.org/10.1007/s13199-019-00637-z>.