



SPECIAL ARTICLE

Mosquito-borne pathogens . . . Now Rickettsias?



Silvina Goenaga ^{a,*}, Estefania Raquel Boaglio^{a,b}, Lucia Victoria Martinez^c,
Juliana Patricia Sanchez^b

^a Instituto Nacional de Enfermedades Virales Humanas (INEVH-ANLIS) "Dr. Julio I. Maiztegui", Pergamino 2700, Argentina

^b Centro de Bioinvestigaciones (CeBio) y Centro de Investigación y Transferencia del Noroeste de Buenos Aires (CITNOBA-CONICET), Universidad Nacional del Noroeste de la Provincia de Buenos Aires, Pergamino 2700, Argentina

^c Laboratorio de Retrovirus y Biología Molecular, Hospital de Pediatría "Prof. Dr. Juan P. Garrahan" Garrahan, CABA, C1245AAM, Argentina

Received 25 October 2024; accepted 14 March 2025

Available online 26 May 2025

KEYWORDS

Rickettsia;
Mosquitoes;
Argentina

Abstract In the last few decades, the emergence of several infectious diseases has increased. Mosquitoes are the most important vectors due to their role in transmitting several pathogens, such as *dengue*, *chikungunya* and *Plasmodium* spp. Recently, mosquitoes have been suspected of transmitting Rickettsias, another emerging pathogen. However, their role in the transmission of this agent remains unclear. In this study, we report the detection of a *Rickettsia* spp. in *Coquillettidia venezuelensis* mosquitoes collected from Corrientes Province, Argentina. The phylogenetic analyses show that this *Rickettsia* is related to *Rickettsia felis*, and the *ompB* sequence shows high homology with *Rickettsia tillamookensis*. The evidence of the detection of *Rickettsia* in mosquitoes shows that several infectious diseases could be transmitted by mosquitoes, and the interaction between them is important to investigate. More studies should be done in order to analyze the impact on public health.

© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Asociación Argentina de Microbiología. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

PALABRAS CLAVE

Rickettsias;
Mosquitos;
Argentina

Patógenos transmitidos por mosquitos. . . ¿Ahora rickettsias?

Resumen En las últimas décadas, la aparición de enfermedades infecciosas ha ido en aumento. Los mosquitos son los vectores más importantes debido a que actúan como transmisores de diversos patógenos, entre ellos, dengue, chikunguña y *Plasmodium* spp. Recientemente, se ha sospechado que los mosquitos también pueden transmitir *rickettsias*, otro patógeno emergente. Sin embargo, el papel de los mosquitos en la transmisión de este agente es aun poco claro. En este estudio informamos la detección de *Rickettsia* spp. en la especie de mosquito *Coquillettidia venezuelensis*, de ejemplares capturados en la provincia de Corrientes, Argentina. Este

* Corresponding author.

E-mail address: sgoenaga@anlis.gob.ar (S. Goenaga).

representa el primer reporte de detección de ADN de *rickettsia* en el país. El análisis filogenético indica que esta *rickettsia* está relacionada con *Rickettsia felis* y la secuencia del gen *ompB* muestra una alta homología con *Rickettsia tillamookensis*. La detección de *rickettsia* en los mosquitos sugiere que estos insectos podrían ser los vectores de múltiples agentes causales de enfermedades infecciosas, lo que resalta la necesidad de investigar la interacción entre ellos. Se deben realizar más estudios con el fin de analizar el impacto de este hallazgo en la salud pública.

© 2025 Los Autores. Publicado por Elsevier España, S.L.U. en nombre de Asociación Argentina de Microbiología. Este es un artículo Open Access bajo la licencia CC BY (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

In recent decades, the impact of several infectious diseases has increased. Their emergence may be attributed to factors such as increasing urbanization, changes in land use, global warming, increased human–animal contact, and the expanded global distribution of their vectors³⁷.

Rickettsiales are a group of bacteria that are obligate intracellular, Gram-negative microorganisms, primarily vectored by arthropods. Their importance in public health lies in the fact that they can cause severe human diseases, such as anaplasmosis, ehrlichiosis, and rickettsioses. Ticks have been considered the most important vectors of *Rickettsia*; however, other ectoparasites, such as fleas, mites, and midges, also act as vectors for different *Rickettsia* species⁷.

Rickettsia is responsible for emerging and re-emerging diseases worldwide³⁰. In nature, *Rickettsia* bacteria are spread by both transstadial and transovarial (i.e., vertical) transmission in their arthropod hosts, or by horizontal (co-feeding) transmission through an infected vertebrate. In Argentina, human cases of *Rickettsia* have been reported, including *R. rickettsii* and *R. parkeri*^{26,27}. Additionally, *R. massiliae* has been detected in ticks⁵, and *R. asembonensis* and *R. felis* have been recorded in fleas^{1,17,20,21,29}.

Mosquitoes are among the most important insects in terms of public health; their blood-feeding behavior is a critical determinant of pathogen transmission. It is well known that mosquitoes are the primary vectors of several arboviruses, such as dengue, chikungunya, and West Nile virus, and are the major vectors of *Plasmodium* spp., the causative agent of malaria³³. Moreover, in recent years, *Rickettsia* have been detected in mosquito samples^{12,18,19,31,32,36}. However, the role of mosquitoes in the transmission of *Rickettsia* and their potential impact on public health remain unclear. Additionally, no studies on *Rickettsia* involving these insects have been conducted in Argentina.

In order to gather more data on *Rickettsia* species in mosquitoes, the objective of this study was to analyze the presence of *Rickettsia* spp. in mosquito samples from north-eastern Argentina using molecular biology methods.

Materials and methods

Mosquito collection and identification

In November 2019, mosquitoes were collected in Tabay, Corrientes (28°18'27"S, 58°17'13"W) during an entomological surveillance for yellow fever virus. This town is located 140 km from the city of Corrientes and has a population of 1700 inhabitants (Censo Nacional de Población, Hogares y Viviendas de la República Argentina, 2010). Manual captures were performed over three consecutive days, during the day, with morning and afternoon sessions lasting four hours (from 8 a.m. to 12 p.m. and from 2 p.m. to 6 p.m., respectively). Mosquitoes were stored in cryovials containing liquid nitrogen and shipped to the Instituto Nacional de Enfermedades Virales Humanas (INEVH-ANLIS) laboratory. Mosquito processing was performed as previously described by Goenaga et al.¹¹. Briefly, upon arrival at the laboratory, the specimens were sorted by species and day of capture under a stereoscopic microscope on a chilled table. Specimens were morphologically identified using a dichotomous key developed by Darsie⁵.

Pools were assembled with mosquitoes of the same species captured on the same day, and each was homogenized in a microtube containing a 3 mm tungsten bead and 1 ml of phosphate-buffered saline solution with 0.75% bovine albumin, penicillin (100 units/ml), and streptomycin (100 µg/ml) for 1 min at 20 cycles per second using a Bead Ruptor 24 Elite (OMNI International, Kennesaw, Georgia, USA). Homogenates were clarified by centrifugation at 5000 × g for 10 min at 4 °C.

Samples were processed using the DNeasy Blood and Tissue Kit (QIAGEN, Hilden, North Rhine-Westphalia, Germany) and eluted into final volumes of 60 µl.

The presence of *Rickettsia* spp. was screened by a specific PCR of the following genes: citrate synthase (*gltA*: 401 bp fragment and 834 bp fragment), outer membrane protein A (*ompA*), and outer membrane protein B (*ompB*) genes (Table 1). For the amplification, the PCR program started with an initial denaturation for 2 min at 98 °C, followed by 40 cycles (94 °C for 15 s, gene-specific annealing at 48 °C for 30 s, and 74 °C for 30 s), and a final extension step at 72 °C for

Table 1 Genes, primer names and sequences, and PCR amplification conditions used for *Rickettsia* detection.

Target gene	Primer name	Nucleotide sequence (5'–3')	Annealing T (°C)	Product length (bp)	Reference
<i>gltA</i>	CS-78	GCAAGTATCGGTGAGGATGTAAT	48	401	15
	CS-323	CTTCCTTAAATCAATAAATCAGGAT			
	CS-239	GCTCTTCTCATCTATGGCTATTAT	48	834	16
<i>ompA</i>	CS-1069	CAGGGTCTTCGTGCATTCTT			
	Rr190.70	ATGGCGAATATTTCTCAAAA	46	530	25
	190-701	GTTCCGTTAATGGCAGCATCT			
<i>ompB</i>	120-M59	CCGCAGGGTTGGTAACTGC	48	856	28
	120-807	CCTTTTAGATTACCGCCTAA			

5 min (Table 1). The PCR reaction was set to a final volume of 20 µL, containing: 25–100 ng of template DNA, 1.5 mM MgCl₂, 0.2 µM of each primer, 0.2 mM of each dNTP, 1X reaction buffer, 0.5 U of *Taq Platinum* DNA polymerase, and ultrapure sterile water to reach the final volume. For each reaction, a negative control (5 µL of molecular-grade water) and a positive control (5 µL of DNA belonging to *R. prowazekii*) were included. PCR products were electrophoresed through a 2% agarose gel, stained with 10 000× SYBR® Safe (Invitrogen, Eugene, Oregon, USA), and examined by LED transillumination.

Conventional PCR products (*gltA*, *ompA*, and *ompB* genes) were purified using the QIAquick Kit (QIAGEN, Hilden, North Rhine-Westphalia, Germany) according to the manufacturer's protocol. Both strands of the purified DNA were sequenced using the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA). Sanger sequencing was performed using an ABI PRISM 3500 DNA Analyzer (Applied Biosystems, Foster City, CA, USA).

Sequences were edited using the BioEdit program¹³. A homology analysis was conducted using the BLAST algorithm (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) against the GenBank nucleotide database to elucidate the identity of each sequence. Finally, the Clustal-W method was used for alignment (available at <http://www.mbio.ncsu.edu/BioEdit/bioedit.html>).

Phylogenetic trees were constructed using the maximum likelihood (ML) method for both the *gltA* and *ompB* genes. For the phylogenetic analysis, the IQ-TREE web server was used³⁵. The best-fit model for *gltA* was TIME+G4, and for *ompB*, TVM+F+G4, both chosen according to the Akaike Information Criterion (AIC)¹⁴. The confidence levels of the nodes were determined using bootstrapping with 1000 replicates. Tree topology was reconstructed using FigTree software²⁴.

Results

We captured 357 individual mosquitoes (Diptera: Culicidae) belonging to the genera *Aedes* (0.28%), *Anopheles* (0.28%), *Coquillettidia* (59.6%), *Mansonia* (15.4%), *Psorophora* (20.7%), *Sabethes* (0.84%), and *Wyeomyia* (2.8%).

Of all the pools (25 in total), only one tested positive for both the *gltA* and *ompB* genes by PCR. No amplification was detected using primers targeting the *ompA* gene. Since the *gltA* gene is highly conserved, its amplification only confirms

the presence of the genus; therefore, to confirm the species identity, the *ompB* gene was sequenced.

A 750 bp fragment corresponding to the *gltA* gene and a 780 bp fragment corresponding to the *ompB* gene were sequenced and deposited in the GenBank nucleotide database under accession numbers PQ488464 (*gltA*) and PQ488463 (*ompB*).

The positive pool was composed of 39 individuals from *Coquillettidia venezuelensis* (Theobald, 1912). For species confirmation, a PCR targeting the cytochrome c oxidase 1 (COI) gene was performed using the primer pairs from Folmer et al.⁸: LCO1490 (5'-GGTCAACAAATCATAAAGATATTGG-3') and HCO2198 (5'-TAAACTTCAGGGTGACCAAAAAATCA-3'), which were used to amplify the ~658 bp fragment of the COI gene. The sequences were deposited in GenBank under accession number PQ315765.


For the *ompB* gene, nBLAST analysis indicated identities of 93.94% (query cover: 100%; e-value: 0) with *Rickettsia tillamookensis* and 91.78% (query cover: 100%; e-value: 0) with *R. felis*, respectively.

In Figure 1, we show the results of the phylogenetic analysis (ML) using a 750 bp fragment from the citrate synthase gene (*gltA*), and Figure 2 shows the analysis using a 780 bp fragment from the *ompB* gene sequences. These analyses demonstrate that the sequences obtained in this study belong to *Rickettsia* (Fig. 1). The *ompB* gene sequence analysis shows that it clusters with *R. felis* and *R. assemblensis*, but not with *R. tillamookensis*, despite the high identity.

Discussion

In this study, we report the presence of *Rickettsia* related to *R. tillamookensis* and *R. felis* in one pool of mosquitoes belonging to *Coquillettidia* (*Rhynchotaenia*) *venezuelensis* (Theobald, 1912), collected in Corrientes province, Argentina. This is the first detection of *Rickettsia* DNA in mosquitoes in Argentina.

The genus *Rickettsia* comprises Gram-negative bacteria that are obligate intracellular pathogens. It currently contains 32 species, divided into five phylogenetic groups: the Spotted Fever group (SFGI and SFGII), with the SFGI *Rickettsia* group containing 24 species (e.g., *R. conorii*, *R. massiliae*, *R. rickettsii*), and the SFGII *Rickettsia* group (also referred to as the transitional group, TRG), which includes five species (e.g., *R. felis* and *R. australis*); the Typhus group (TG), which includes *R. prowazekii* and






Figure 2 Phylogenetic tree generated using the maximum likelihood method for *Rickettsia* spp., based on the *ompB* gene. The strain of interest is identified as 7244 (GenBank accession number PQ488463), with bootstrap values indicated at the nodes.

associated with this agent²³. However, in Argentina, despite several reports of *R. felis* detection, there are no reports in patients, and it is possible that the prevalence of the infection is underestimated. Healthcare professionals should be alerted to the possibility of infection with *R. felis* among their patients. The first detection of *R. felis* in mosquitoes was made by Socolovschi et al.³¹. Following that, Guo et al.¹² reported the presence of *Rickettsia* species (*R. japonica*, *R. sibirica*, and *R. monacensis*) in adult mosquitoes using groEL sequences. In addition, *R. japonica*, *R. sibirica*, and *R. monacensis* were detected in all life stages (egg, larva, pupa, and adult). These data show the presence of several *Rickettsia* species; however, there is no evidence of the role mosquitoes play in transmission.

A novel *Rickettsia* species was detected in *Anopheles* mosquitoes in China, showing high similarity to *R. felis*¹⁸. In the phylogenetic analysis, our strain did not align with this new *Rickettsia*. This result suggests that it could represent a new *Rickettsia* species also related to *R. felis*, but further studies are needed to confirm this hypothesis.

In order to determine the role of mosquitoes as vectors, one approach should be to isolate the bacterium from the saliva or salivary glands of specimens from wild-caught mosquitoes. Additionally, the infection of mosquitoes following experimental feeding on a bacteremic host, as well

as the transmission of bacteria to a vertebrate through the bite of a mosquito, has been demonstrated. Dieme et al.⁶ infected *Anopheles gambiae* mosquitoes orally with *R. felis* and reported the transmission potential of *R. felis* 15 days post-infection, both via salivation and by detecting *R. felis* in mice after feeding on mosquitoes that had been previously infected. However, there was no report of vertical transmission, which suggests that horizontal transmission occurs in nature.

Overall, the studies indicate that mosquitoes may play a role in the transmission of several species of *Rickettsia*. However, the epidemiological significance of this finding remains uncertain. The detection of *Rickettsia* in mosquitoes suggests that they could transmit various infectious diseases, making it crucial to analyze their interactions. Further research is needed to assess the impact on public health and the potential risk of a new emerging zoonotic disease.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT to improve certain sentences in English. After using this

tool, the authors reviewed and edited the content as needed and took full responsibility for the publication's content.

Funding

This research was funded by INEVH (ANLIS) and funded partially by Subsidios de Investigación Biauiales (SIB) 2022 EXP-2164/2022 UNNOBA.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgment

The authors would like to express their gratitude to Carina Sen for assistance in bioinformatics analysis.

References

- Acosta DB, Zanocco FA, Ruiz M, Sánchez JP. Domestic dogs as hosts of ectoparasites carrying *Rickettsia*, *Bartonella*, and *Mycoplasma* in urban, peri-urban, and rural areas from central Argentina. *Mastozool Neotrop*. 2023;30:1–12.
- Aitken TH, Downs WG, Anderson CR, Spence L, Casals J. Mayaro virus isolated from a Trinidadian mosquito, *Mansonia venezuelensis*. *Science*. 1960;131:986, <http://dx.doi.org/10.1126/science.131.3405.986>.
- Anderson CR, Spende L, Downs WG, Aitken TH. Oropouche virus: a new human disease agent from Trinidad, West Indies. *Am J Trop Med Hyg*. 1961;10:574–8, <http://dx.doi.org/10.4269/ajtmh.1961.10.574>.
- Campos RE, Laurito M, Muttis E. *Culicidae* (Diptera) species from Argentina and Uruguay; 2024. Retrieved from: <https://biodar.unlp.edu.ar/culicidae/>
- Darsie RF. Mosquitoes of Argentina. Part I. Keys for identification of adult females and fourth stages larvae in English and Spanish (*Diptera: Culicidae*). *Mosquito Syst*. 1985;17:153–253.
- Dieme C, Bechah Y, Socolovschi C, Audoly G, Berenger JM, Faye O, Raoult D, Parola P. Transmission potential of *Rickettsia felis* infection by *Anopheles gambiae* mosquitoes. *Proc Natl Acad Sci U S A*. 2015;112:8088–93, <http://dx.doi.org/10.1073/pnas.1413835112>.
- El Karkouri K, Ghigo E, Raoult D, Fournier PE. Genomic evolution and adaptation of arthropod-associated *Rickettsia*. *Sci Rep*. 2022;12:3807, <http://dx.doi.org/10.1038/s41598-022-07725-z>.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol*. 1994;3:294–9 [PMID: 7881515].
- Forattini OP. *Entomologia médica*. Culicini: Haemagogus, Mansonia, Culiseta, Sabethini, Toxorhynchitini, Arboviroses, Filariose bancroftiana, Genética, vol. 3. São Paulo: Editora da USP; 1965.
- Gauthier DT, Karpathy SE, Grizzard SL, Batra D, Rowe LA, Paddock CD. Characterization of a novel transitional group *Rickettsia* species (*Rickettsia tillamookensis* sp. nov.) from the western black-legged tick, *Ixodes pacificus*. *Int J Syst Evol Microbiol*. 2021;71:004880, <http://dx.doi.org/10.1099/ijsem.0.004880>.
- Goenaga S, Fabbri CM, García JB, Rondán JC, Gardenal N, Calderón GE, Enría DA, Levis SM. New strains of *Culex flavivirus* isolated in Argentina. *J Med Entomol*. 2014;51:900–6, <http://dx.doi.org/10.1603/me13172>.
- Guo WP, Tian JH, Lin XD, Ni XB, Chen XP, Liao Y, Yang SY, Dummer JS, Holmes EC, Zhang YZ. Extensive genetic diversity of *Rickettsiales* bacteria in multiple mosquito species. *Sci Rep*. 2016;6:38770, <http://dx.doi.org/10.1038/srep38770>.
- Hall T. BioEdit 6.0.7. Department of Microbiology, North Carolina State University; 2004. Retrieved from: <http://www.mbio.ncsu.edu/BioEdit/bioedit.html>
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermini LS. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*. 2017;14:587–9, <http://dx.doi.org/10.1038/nmeth.4285>.
- Labruna MB, Whitworth T, Horta MC, Bouyer DH, McBride JW, Pinter A, Popov V, Gennari SM, Walker DH. *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the state of São Paulo, Brazil, where Brazilian spotted fever is endemic. *J Clin Microbiol*. 2004;42:90–8, <http://dx.doi.org/10.1128/JCM.42.1.90-98.2004>.
- Labruna MB, McBride JW, Bouyer DH, Camargo LM, Camargo EP, Walker DH. Molecular evidence for a spotted fever group *Rickettsia* species in the tick *Amblyomma longirostre* in Brazil. *J Med Entomol*. 2004;41:533–7, <http://dx.doi.org/10.1603/0022-2585-41.3.533>.
- López Berrizbeitia MF, Acosta DB, Sánchez JP. Wild rodent fleas carrying *Bartonella* and *Rickettsia* in an area endemic for vector-borne diseases from Argentina. *Sci Rep*. 2024;14:23269, <http://dx.doi.org/10.1038/s41598-024-74786-7>.
- Lu M, Chen S, Meng C, Wang W, Li H, Sun Y, Li M, Ma X, Ma Y, Duan C, Li K. A novel *Rickettsia* species closely related to *Rickettsia felis* in *Anopheles* mosquitoes from Yingkou City, Northeast China. *Zoonoses Public Health*. 2023;70:568–71, <http://dx.doi.org/10.1111/zph.13043>.
- Mediannikov O, Socolovschi C, Edouard S, Fenollar F, Mouffok N, Bassene H, Diatta G, Tall A, Niangaly H, Doumbo O, Lekana-Douki JB, Znazen A, Sari H, Ratmanov P, Richet H, Ndiath MO, Sokhna C, Parola P, Raoult D. Common epidemiology of *Rickettsia felis* infection and malaria, Africa. *Emerg Infect Dis*. 2013;19:1775–83, <http://dx.doi.org/10.3201/eid1911.130361>.
- Melis M, Espinoza-Carniglia M, Savchenko E, Nava S, Lareschi M. Molecular detection and identification of *Rickettsia felis* in *Polygenis* (Siphonaptera, Rhopalopsyllidae, Rhopalopsyllinae) associated with cricetid rodents in a rural area from central Argentina. *Vet Parasitol Reg Stud Rep*. 2020;21:100445, <http://dx.doi.org/10.1016/j.vprsr.2020.100445>.
- Nava S, Pérez-Martínez L, Venzal JM, Portillo A, Santibañez S, Oteo JA. *Rickettsia felis* in *Ctenocephalides felis* from Argentina. *Vector Borne Zoonotic Dis*. 2008;8:465–6, <http://dx.doi.org/10.1089/vbz.2007.0243>.
- Paddock CD, Slater K, Wei A, Zambrano ML, Kleinjan JE, Padgett KA, Saunders MEM, Andrews ES, Trent E, Zhong J, Sambado S, Goldsmith CS, Pascoe EL, Foley J, Lane RS, Karpathy SE. Detection and isolation of *Rickettsia tillamookensis* (*Rickettsiales: Rickettsiaceae*) from *Ixodes pacificus* (Acari: Ixodidae) from multiple regions of California. *J Med Entomol*. 2022;59:1404–12, <http://dx.doi.org/10.1093/jme/tjac038>.
- Parola P. *Rickettsia felis*: from a rare disease in the USA to a common cause of fever in sub-Saharan Africa. *Clin Microbiol Infect*. 2011;17:996–1000, <http://dx.doi.org/10.1111/j.1469-0691.2011.03516.x>.
- Rambaut A. Figtree, a graphical viewer of phylogenetic trees; 2014. Retrieved from: <http://treebioedacuk/software/figtree>
- Regnery RL, Spruill CL, Plikaytis BD. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol*. 1991;173:1576–89, <http://dx.doi.org/10.1128/jb.173.5.1576-1589.1991>.
- Ripoll CM, Remondégui CE, Ordóñez G, Arazamendi R, Fusaro H, Hyman MJ, Paddock CD, Zaki SR, Olson JG, Santos-Buch CA. Evidence of rickettsial spotted fever

- and ehrlichial infections in a subtropical territory of Jujuy, Argentina. *Am J Trop Med Hyg.* 1999;61:350–4, <http://dx.doi.org/10.4269/ajtmh.1999.61.350>.
27. Romer Y, Seijo AC, Crudo F, Nicholson WL, Varela-Stokes A, Lash RR, Paddock CD. *Rickettsia parkeri* Rickettsiosis, Argentina. *Emerg Infect Dis.* 2011;17, <http://dx.doi.org/10.3201/eid1707.101857>.
28. Roux V, Raoult D. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (ompB). *Int J Syst Evol Microbiol.* 2000;50:1449–55, <http://dx.doi.org/10.1099/00207713-50-4-1449>.
29. Ruiz M, Acosta DB, Baricalla A, Sánchez JP. Molecular detection of *Rickettsia* in ectoparasites (*Siphonaptera* and *Phthiraptera*) of domestic and feral pigs from Argentina. *Parasitol Res.* 2021;120:3611–8, <http://dx.doi.org/10.1007/s00436-021-07291-9>.
30. Rymaszewska A, Piotrowski M. *Rickettsia* species: genetic variability vectors rickettsiosis – a review. *Pathogens.* 2024;13:661, <http://dx.doi.org/10.3390/pathogens13080661>.
31. Socolovschi C, Pagés F, Ndiath MO, Ratmanov P, Raoult D. *Rickettsia* species in African *Anopheles* mosquitoes. *PLoS One.* 2012;7:e48254, <http://dx.doi.org/10.1371/journal.pone.0048254>.
32. Socolovschi C, Pagés F, Raoult D. *Rickettsia felis* in *Aedes albopictus* mosquitoes, Libreville, Gabon. *Emerg Infect Dis.* 2012;18:1687–9, <http://dx.doi.org/10.3201/eid1810.120178>.
33. Soni S, Gill VJS, Anusheel, Singh J, Chhabra J, Gill GJS, Bakshi R. Dengue, Chikungunya, and Zika: the causes and threats of emerging and re-emerging arboviral diseases. *Cureus.* 2023;15:e41717, <http://dx.doi.org/10.7759/cureus.41717>.
34. Stein M, Zalazar L, Willener JA, Almeida FL, Almirón WR. *Culicidae* (Diptera) selection of humans, chickens and rabbits in three different environments in the province of Chaco, Argentina. *Mem Inst Oswaldo Cruz.* 2013;108:563–71, <http://dx.doi.org/10.1590/s0074-02762013000500005>.
35. Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 2016;44:W232–5, <http://dx.doi.org/10.1093/nar/gkw256>.
36. Zhang J, Lu G, Li J, Kelly P, Li M, Wang J, Huang K, Qiu H, You J, Zhang R, Wang Y, Zhang Y, Wu H, Wang C. Molecular detection of *Rickettsia felis* and *Rickettsia bellii* in mosquitoes. *Vector Borne Zoonotic Dis.* 2019;19:802–9, <http://dx.doi.org/10.1089/vbz.2019.2456>.
37. Zhou S, Liu B, Han Y, Wang Y, Chen L, Wú Z, Yang J. ZOVER: the database of zoonotic and vector-borne viruses. *Nucleic Acids Res.* 2021;50:D943–9, <http://dx.doi.org/10.1093/nar/gkab862>.