



ORIGINAL ARTICLE

Microbial consortia as a biocontrol strategy for bacterial blight in basmati rice



Bestha Sai Teja^a, Vishal Gupta^a, Sushil Kumar Gupta^b, Ayushi Sharma^a,
Zakir Amin^{Id}^c, Fayaz Ahmad Mohiddin^d, Suhail Ashraf^e, Mona Saleh Al Tami^{f,*},
Khalid E. Hamed^g

^a Division of Plant Pathology, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Chatha 180009, India

^b Directorate of Research, Sher-e-Kashmir University of Agricultural Sciences & Technology of Jammu, Chatha 180009, India

^c Division of Plant Pathology, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, Wadura 193201, India

^d Section of Plant Pathology, Mountain Research Centre for Field Crops (MRCFC), Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Khudwani, Kashmir 192101, India

^e Department of Plant Biotechnology, Center for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, 641003 Coimbatore, Tamil Nadu, India

^f Department of Biology, College of Science, Qassim University, PO Box 6622, Buraidah 51452, Qassim, Saudi Arabia

^g Department of Plant Protection, College of Agriculture and Food, Qassim University, PO Box 6622, Buraidah 51452, Qassim, Saudi Arabia

Received 22 January 2025; accepted 28 August 2025

Available online 8 October 2025

KEYWORDS

Basmati rice;
Bacterial leaf blight;
Microbial consortium;
Yield

Abstract Basmati rice is in high demand in the world market because of its fine, soft, long and distinctive aromatic grains. Bacterial leaf blight (BLB), caused by *Xanthomonas oryzae* pv. *oryzae*, is a destructive disease, resulting in enormous yield loss in basmati rice. Chemical management against BLB is effective but not the preferred approach because of its detrimental effects on humans, environment and ecological disruption. In the present study, seed treatment, seedling treatment and foliar application with microbial antagonists, such as *Trichoderma harzianum* and *Pseudomonas fluorescens*, and a consortium encompassing both microorganisms plus *Trichoderma asperellum*, were assessed against BLB in basmati rice. Among all the bioformulation treatments, seed treatment with the microbial consortium revealed significant ($p < 0.05$) results, achieving maximum inhibition of BLB incidence, intensity and lesion length of 48.56, 52.42 and 45.41 percent respectively, compared to the untreated control. Simultaneously, the same treatment also resulted in significant ($p < 0.05$) enhancement of growth attributes, such as panicle length (55.83%), plant fresh weight (15.55%), grains per panicle (38.23%), test weight (55.49%) and yield (47.68%). A maximum increase in superoxide dismutase and hydrogen peroxide activities by 81.58 and 84.29 percent respectively over the control was

* Corresponding author.

E-mail address: Tamie@qu.edu.sa (M.S. Al Tami).

also recorded. The results of the study suggest that the effectiveness of BLB management under field conditions applying the microbial consortium (*T. harzianum* + *T. asperellum* + *P. fluorescens*) was similar to that of the chemical control. Further, efforts are being made to evaluate the bio-control activity of the microbial consortium in large-scale field experiments and its commercial potential.

© 2025 The Author(s). Published by Elsevier España, S.L.U. on behalf of Asociación Argentina de Microbiología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Arroz basmati;
Tizón bacteriano de
las hojas;
Consortios
microbianos;
Rendimiento

Consortios microbianos como estrategia de biocontrol para el tizón bacteriano en el arroz basmati

Resumen El arroz basmati es sumamente valorado en el mercado global por sus granos largos, finos, suaves y aromáticos. El tizón bacteriano de las hojas (BLB), causado por *Xanthomonas oryzae* pv. *oryzae*, representa una seria amenaza en el cultivo del arroz basmati, generando pérdidas significativas en el rendimiento. Aunque el manejo químico de este patógeno ha demostrado eficacia, esta no es una práctica sostenible debido a sus efectos negativos sobre la salud humana, el ambiente y el equilibrio ecológico. En este estudio se evaluaron distintos antagonistas microbianos (*Trichoderma harzianum*, *Pseudomonas fluorescens* y un consorcio que incluyó dichos microorganismos más *Trichoderma asperellum*) y diferentes métodos de aplicación (tratamiento de semillas, de plántulas y aplicación foliar) para el manejo del BLB en el cultivo de arroz basmati. El tratamiento con el consorcio microbiano aplicado en semillas fue el más eficaz, logrando reducciones significativas ($p < 0,05$) en la incidencia (48,56%), la intensidad (52,42%) y la longitud de la lesión (45,41%) respecto al control. Este tratamiento también promovió un aumento notable en indicadores agronómicos como la longitud de la panícula (55,83%), el peso fresco de la planta (15,55%), el número de granos por panícula (38,23%), el peso de la prueba (55,49%) y el rendimiento (47,68%). Además, se observaron incrementos en la actividad de superóxido dismutasa (81,58%) y peróxido de hidrógeno (84,29%), lo que indica una respuesta de defensa mejorada. Se concluye que el consorcio microbiano evaluado representa una alternativa eficaz y sostenible para reducir el impacto del BLB en el cultivo de arroz basmati. Se están llevando a cabo estudios para evaluar su aplicación a gran escala y el potencial comercial.

© 2025 Los Autores. Publicado por Elsevier España, S.L.U. en nombre de Asociación Argentina de Microbiología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Basmati is an extraordinarily premium, long-grain rice variety characterized by a pleasant aroma³². Because of its fine, soft and nutrient-rich grains, basmati is highly demanded in international, national and regional markets. India is the leading producer of basmati rice, with 74 percent of the world's supply, followed by Pakistan, the Philippines and China²⁷. Basmati rice is also well known for its health benefits, including good bioavailability of cystine, vitamin B1 and melatonin, its low glycemic index and its role in fighting diabetes². However, the productivity of basmati rice is threatened due to the outbreak of several biotic stresses, such as bacterial leaf blight, blast and bakanae disease²².

Bacterial leaf blight (BLB), caused by *Xanthomonas oryzae* pv. *oryzae* (*Xoo*), is one of the major diseases affecting basmati rice. It is responsible for 20–80 percent of yield loss in India⁴¹, 30 percent in Japan⁴⁴, 50 percent in Malaysia³⁵, 80 percent in Indonesia⁴³, 50 percent in the Philippines³, 20–40 percent in Bangladesh¹⁶ and up to 80

percent in Mali⁴⁵. Diverse pathogenic variations among the *Xoo* populations have been reported in the basmati-growing regions of Jammu and Kashmir⁶. *Xoo* penetrates the host through hydathodes and wounds present on the surface of the plant, eventually affecting the parenchymatous cells of xylem, traveling vertically and clogging the xylem vessels. Subsequently, bacterial ooze is observed coming out from the hydathodes²¹. *Xoo* disease-causing ability is dependent on the type-3 secretion system (T3SS), which is important for infection of susceptible hosts and plant defense activation¹⁵.

Management of bacterial leaf blight can be achieved through chemical means, but the overuse of the antibiotic tetracycline causes several allergic and metabolic disorders in humans, potentially resulting in anti-microbial resistance⁴. The alternative to chemical pesticides is the biological management of plant diseases, which is efficient and effective through the production of antibiotics, hydrogen cyanide, siderophores, lytic enzymes and antimicrobial compounds against phytopathogens. Moreover, biocontrol improves the growth and development of the host plant³⁴

by unveiling bio-stimulant traits, such as increasing the bioavailability and solubilization of various soil nutrients for plant growth, stimulating phytohormones such as auxins, cytokinins and gibberellins, and producing metabolites that improve tolerance to abiotic stress⁹. Biocontrol agents such as *Trichoderma* spp., *Pseudomonas* spp. and *Bacillus* spp. are known to exhibit several bioinsecticidal, bionematicidal and biopesticidal activities¹², and are used individually or in combination as a consortium.

A microbial consortium is a combination of multiple strains or species of beneficial microbes gaining more importance than a single inoculant because of their multifaceted mechanisms of disease mitigation, enhancement of soil properties and improvement of growth and yield parameters of the host plant²⁵. Further, the occasional inconsistent, ineffective and variable performance of single inoculants influenced by varying environmental conditions limit their competitiveness and effectiveness against plant pathogens⁴⁶. The application of synthetic microbial communities (SynComs) to improve crop productivity by mitigating plant diseases, improving soil and plant health is a major area of research in this decade⁸. Therefore, this investigation was aimed at: (i) evaluating the microbial consortium against bacterial leaf blight in basmati rice and (ii) studying the effect of the consortium on plant growth and induced systemic resistance (ISR).

Materials and methods

Experimental details

The seed of Basmati-370 used in the experiment was obtained from the Mega Seed Unit of SKUAST-Jammu, Chatha. The biocontrol formulation of each inoculant along with the microbial consortium consisting of *Trichoderma harzianum* (GenBank accession no. MN721820), *Trichoderma asperellum* (GenBank accession no. MT395682), and *Pseudomonas fluorescens* (GenBank accession no. KJ194131), along with the pathogen *X. oryzae* pv. *oryzae* (GenBank accession no. MN718665) were obtained from the Biocontrol Laboratory, Advanced Centre for Horticulture Research, SKUAST-Jammu, Udhewala.

Artificial inoculation of *X. oryzae* pv. *oryzae*

X. oryzae pv. *oryzae* (Xoo) was mass-multiplied in a nutrient broth medium (Hi-media) and artificially inoculated onto the paddy crop using the leaf clipping method¹⁸. The top portion of fully developed leaves of Basmati-370, measuring between 2.5 and 6 cm, were clipped off using sterilized scissors that had already been dipped in a suspension of Xoo inoculum (1×10^8 CFU/ml) during the tillering stage.

Effectiveness of the microbial consortium against bacterial leaf blight of rice under field conditions

Experimental plots were set up in the Research Field of the Division of Plant Pathology (32°39.1749' N; 74°48.4268' E) during the Kharif season of 2023–2024. Rice variety Basmati-370 was planted in plots (3 × 4 m²) arranged in a

randomized block design (RBD) with three replications. Each plot had rows spaced 20 cm apart and plants spaced 15 cm apart. Eleven treatments were applied to manage bacterial leaf blight of rice, whose details are mentioned in Table 1. Prior to sowing, the seeds were surface-sterilized with 0.1 percent of sodium hypochlorite. After sterilization, seeds were soaked separately in each inoculant and the microbial consortium consisting of *T. harzianum*, *P. fluorescens* and *T. asperellum*, then shade-dried and sown in plots on 24th June, 2023. Additionally, seedling treatment was carried out by soaking 30 day-old seedlings of untreated plants for 20 min in their respective treatments consisting of single inoculants and the microbial consortium. Similarly, foliar application of the respective bioagents and streptocycline + copper oxychloride was carried out using a knapsack sprayer pump after the 7th day of pathogen inoculation.

Biochemical parameters

Fresh leaf samples were collected from each plot after seven days of pathogen inoculation and assessed for the quantification of superoxide dismutase and hydrogen peroxide (H₂O₂)⁴².

Observations recorded

The disease attributes such as percent incidence, percent intensity,³⁸ lesion length (cm) of BLB were recorded after 14 days of Xoo inoculation. Growth attributes such as fresh and dry root weight (g) were recorded at 30, 60 and 90 days after transplant (DAT), and panicle length (cm), plant fresh weight (g), grains per panicle, test weight (g), and yield (q/ha) were recorded at the time of harvesting.

Statistical analysis

Data were analyzed using ANOVA with Duncan multiple range tests in R software, and the correlation analysis was performed using the psych package of R software⁴⁸. The graphical plots were made using Originlab software¹⁰. Additionally, the correlation analysis was performed among growth attributes (panicle length, plant fresh weight, fresh and dry root weight at 30, 60 and 90 DAT, number of grains per panicle, test weight and yield), biochemical attributes (SOD and H₂O₂) and disease attributes (percent intensity, incidence and lesion length) to determine the level of association of each attribute with the incidence of bacterial leaf blight.

Results

Effect of the microbial consortium on the incidence, intensity and lesion length of bacterial leaf blight of rice under field conditions

The effectiveness of the microbial consortium (*T. harzianum* + *T. asperellum* + *P. fluorescens*), single inoculants (*T. harzianum* and *P. fluorescens*) and streptocycline + copper oxychloride were assessed against the bacterial leaf blight caused by *X. oryzae* pv. *oryzae* (Xoo).

Table 1 Different treatments tested for the management of bacterial leaf blight in rice.

Treatment	Details
T ₁	Seed treatment with <i>Trichoderma harzianum</i> @ 1×10^7 spores/ml
T ₂	Seed treatment with <i>Pseudomonas fluorescens</i> @ 1×10^8 CFU/ml
T ₃	Seed treatment using the microbial consortium [<i>T. harzianum</i> (1×10^7 spores/ml) + <i>T. asperellum</i> (1×10^7 spores/ml) + <i>P. fluorescens</i> (2×10^8 CFU/ml)]
T ₄	Seedling treatment with <i>Trichoderma harzianum</i> @ (1×10^7 spores/ml)
T ₅	Seedling treatment with <i>Pseudomonas fluorescens</i> @ (1×10^8 CFU/ml)
T ₆	Seedling treatment with the microbial consortium [<i>T. harzianum</i> (1×10^7 spores/ml) + <i>T. asperellum</i> (1×10^7 spores/ml) + <i>P. fluorescens</i> (2×10^8 CFU/ml)]
T ₇	Foliar application with <i>Trichoderma harzianum</i> at 0.1 percent (1×10^7 spores/ml)
T ₈	Foliar application with <i>Pseudomonas fluorescens</i> at 0.1 percent (1×10^8 CFU/ml)
T ₉	Foliar application using the microbial consortium at 0.1 percent [<i>T. harzianum</i> (1×10^7 spores/ml) + <i>T. asperellum</i> (1×10^7 spores/ml) + <i>P. fluorescens</i> (2×10^8 CFU/ml)]
T ₁₀	Foliar application with streptocycline (40 g) + copper oxychloride (200 g)
T ₁₁	Control (untreated)

Table 2 Effect of the microbial consortium on percent disease incidence, intensity and lesion length of bacterial leaf blight in rice.

Treatment	Percent incidence*	Percent intensity*	Lesion length* (cm)
T ₁	31.43 ± 2.63 ^e	27.43 ± 2.63 ^e	33.46 ± 0.85 ^e
T ₂	34.10 ± 1.60 ^{de}	30.10 ± 1.61 ^{de}	35.38 ± 1.28 ^e
T ₃	27.90 ± 2.84 ^f	23.90 ± 2.85 ^f	27.93 ± 0.70 ^f
T ₄	36.46 ± 1.49 ^{cd}	32.47 ± 1.50 ^{cd}	39.16 ± 0.95 ^d
T ₅	39.40 ± 2.30 ^c	35.40 ± 2.31 ^c	40.93 ± 1.20 ^{cd}
T ₆	28.13 ± 1.00 ^f	24.13 ± 1.01 ^f	30.26 ± 0.80 ^f
T ₇	45.73 ± 1.48 ^b	41.73 ± 1.48 ^b	43.30 ± 0.85 ^{bc}
T ₈	47.20 ± 1.11 ^b	43.20 ± 1.11 ^b	45.76 ± 1.56 ^b
T ₉	45.76 ± 1.90 ^b	41.77 ± 1.90 ^b	44.20 ± 0.90 ^b
T ₁₀	24.40 ± 0.91 ^g	20.40 ± 0.92 ^g	24.46 ± 0.85 ^g
T ₁₁	54.23 ± 3.01 ^a	50.23 ± 3.01 ^a	51.17 ± 4.35 ^a

Different alphabetical letters on the superscript within the columns indicate significant differences among the treatments according to Duncan's multiple range test at $p \leq 0.05$.

T₁ to T₁₁ in the table correspond to the treatment details mentioned in Table 1.

* Data are the average of three replications, ± represents standard deviations of the mean.

Streptocycline + copper oxychloride exhibited minimum percent disease incidence, percent intensity and lesion length (cm) of 24.40 ± 0.91 , 20.40 ± 0.92 and 24.46 ± 0.85 , respectively, followed by seed treatment with the microbial consortium (*T. harzianum* + *T. asperellum* + *P. fluorescens*) resulting in 27.90 ± 2.84 , 23.90 ± 2.85 and 27.93 ± 0.70 cm, respectively. Maximum percent incidence of 54.23 ± 3.01 , percent intensity of 50.23 ± 3.01 and lesion length of 51.23 ± 4.35 cm was recorded in the control (Table 2 and Fig. 1A).

Effect of the microbial consortium on fresh and dry root weight of rice plants

The maximum enhancement of fresh root weight was exhibited in the seed treatment involving the microbial consortium containing *T. harzianum* + *T. asperellum* + *P. fluorescens*, with values of 22.20 ± 0.81 , 25.60 ± 0.81 and 28 ± 0.81 g at 30, 60 and 90 days after transplanting

(DAT), respectively, followed by foliar treatment with streptocycline + copper oxychloride exhibiting 21 ± 0.55 , 24.40 ± 0.55 and 26.80 ± 0.55 g at 30, 60 and 90 DAT, respectively. Additionally, minimum fresh root weight was observed in the control (12.160 ± 0.80 , 15.56 ± 0.80 , 17.96 ± 0.80 g, respectively). Seed treatment using the microbial consortium resulted in maximum dry root weight of 15.20 ± 0.81 , 18.60 ± 0.81 and 21.00 ± 0.81 g after 30, 60 and 90 DAT respectively, followed by foliar application of streptocycline + copper oxychloride (14 ± 0.55 , 17.40 ± 0.55 and 19.80 ± 0.55 g, respectively). Minimum dry root weight (7.80 ± 0.45 , 11.20 ± 1.47 and 13.60 ± 0.85 g, respectively) was observed in the control (Table 3 and Fig. 1B).

Effect of the microbial consortium on plant growth promotion and yield attributes of rice plants

Among all the treatments, foliar application of streptocycline + copper oxychloride showed maximum enhancement




Figure 1 Effects of the microbial consortium on rice. (A) Percentage reduction in the incidence, intensity, and lesion length of bacterial leaf blight; (B) percentage increase in fresh and dry root weight after 30, 60, and 90 days of transplanting; (C) percentage increase in panicle length, fresh weight, grains per panicle, test weight, and yield; (D) superoxide dismutase activity (U/ml) and hydrogen peroxide levels (nMol H_2O_2 /g FW).

of panicle length, plant fresh weight, grains per panicle, test weight and yield with 30.06 ± 2.95 g, 197.10 ± 2.95 g, 74.62 ± 3.16 , 26.66 ± 0.51 g and 22.16 ± 0.66 q/ha, respectively, followed by seed treatment with microbial consortium, exhibiting 25.43 ± 0.61 cm, 194.36 ± 4.53 g, 67.66 ± 4.12 , 22.76 ± 1.10 g and 42.36 ± 0.23 q/ha, respectively. Furthermore, in the control, a panicle length of only 11.23 ± 0.14 cm, plant fresh weight of 164.13 ± 4.10 g, grains per panicle of 41.80 ± 5.14 , test weight of 10.13 ± 0.35 g and yield of 22.16 ± 0.66 q/ha were recorded (Table 4 and Fig. 1C).

Effect of the microbial consortium on superoxide dismutase and hydrogen peroxide in rice plants

The seed treatment with the microbial consortium containing *T. harzianum* + *T. asperellum* + *P. fluorescens*, resulted in maximum superoxide dismutase and hydrogen peroxide with 0.723 ± 0.01 U/ml and 0.705 ± 0.01 nMol H_2O_2 /g FW, respectively, followed by seedling treatment using the microbial consortium (0.631 ± 0.02 U/ml and 0.621 ± 0.02 nMol H_2O_2 /g FW, respectively). Minimum activity of superoxide and hydrogen peroxide was recorded in the control with values of 0.113 ± 0.006 U/ml and 0.130 ± 0.006 nMol H_2O_2 /g FW, respectively (Table 5 and Fig. 1D).

Correlation analysis of percent BLB incidence, intensity, lesion length, fresh and dry root weight, and growth and yield attributes in rice plants

The correlation analysis of several factors, such as percent disease intensity, fresh and dry root weights (30, 60 and 90 DAT), yield, test weight, panicle length, plant fresh weight and lesion length of bacterial leaf blight was conducted along with disease incidence, to determine the association between them. BLB intensity and disease lesion length were positively correlated with disease incidence, with a correlation coefficient (R) of 0.99 and 0.99, respectively. Moreover, fresh and dry root weight (30, 60, 90 DAT), yield, test weight, panicle length and plant fresh weight had a significantly negative correlation with disease incidence, with correlation coefficients (R) of -0.96 , -0.96 , -0.96 , -0.95 , -0.95 , -0.95 , -0.98 , -0.93 , -0.98 , -0.93 and -0.91 , respectively (Supplementary Figure 1).

Correlation analysis of superoxide dismutase (SOD) and hydrogen peroxide (H_2O_2) with percent incidence of BLB in rice plants

Biochemical factors, viz., superoxide dismutase (SOD) and hydrogen peroxide (H_2O_2) showed a significantly but moderately negative correlation with the incidence of BLB, with

Table 3 Effect of the microbial consortium on fresh and dry root weight of rice plants after 30, 60 and 90 days of transplanting.

Treatment	Fresh root weight* (g)			Dry root weight* (g)		
	30 DAT**	60 DAT**	90 DAT**	30 DAT**	60 DAT**	90 DAT**
T ₁	19.00 ± 0.40 ^d	22.40 ± 0.40 ^d	24.80 ± 0.40 ^d	12.00 ± 0.40 ^d	15.40 ± 0.40 ^d	17.80 ± 0.40 ^d
T ₂	18.90 ± 0.50 ^d	22.30 ± 0.05 ^d	24.70 ± 0.50 ^d	11.90 ± 0.50 ^d	15.30 ± 0.50 ^d	17.70 ± 0.50 ^d
T ₃	22.20 ± 0.81 ^a	25.60 ± 0.81 ^a	28.00 ± 0.81 ^a	15.20 ± 0.81 ^a	18.60 ± 0.81 ^a	21.00 ± 0.81 ^a
T ₄	17.60 ± 0.40 ^e	21.00 ± 0.40 ^e	23.40 ± 0.40 ^e	10.60 ± 0.40 ^e	14.00 ± 0.40 ^e	16.40 ± 0.40 ^e
T ₅	17.40 ± 0.50 ^e	20.80 ± 0.50 ^e	23.20 ± 0.50 ^e	10.40 ± 0.50 ^e	13.80 ± 0.50 ^{de}	16.20 ± 0.50 ^e
T ₆	20.00 ± 0.62 ^c	23.40 ± 0.62 ^c	25.80 ± 0.62 ^c	13.00 ± 0.62 ^c	16.40 ± 0.62 ^{bc}	18.80 ± 0.62 ^c
T ₇	16.10 ± 0.36 ^f	19.50 ± 0.36 ^f	21.90 ± 0.36 ^f	9.10 ± 0.36 ^f	12.50 ± 0.36 ^f	14.90 ± 0.36 ^f
T ₈	15.96 ± 0.50 ^f	19.36 ± 0.50 ^f	21.76 ± 0.50 ^f	8.96 ± 0.50 ^f	12.36 ± 0.50 ^f	14.76 ± 0.50 ^f
T ₉	16.30 ± 0.36 ^f	19.70 ± 0.36 ^f	22.10 ± 0.36 ^f	9.30 ± 0.36 ^f	12.70 ± 0.36 ^{ef}	15.10 ± 0.36 ^f
T ₁₀	21.00 ± 0.55 ^b	24.40 ± 0.55 ^b	26.80 ± 0.55 ^b	14.00 ± 0.55 ^b	17.40 ± 0.55 ^b	19.80 ± 0.55 ^b
T ₁₁	12.16 ± 0.80 ^g	15.56 ± 0.80 ^g	17.96 ± 0.80 ^g	7.80 ± 0.45 ^g	11.20 ± 1.47 ^g	13.60 ± 0.85 ^g

Different alphabetical letters on the superscript within the columns indicate significant differences among the treatments according to Duncan's multiple range test at $p \leq 0.05$.

T₁ to T₁₁ in table denote the treatment details mentioned in Table 1.

* Data are the average of three replications, $\pm$ represents standard deviations of the mean.

** Days after transplant.

Table 4 Effect of the microbial consortium on plant growth promotion and yield attributes of rice.

Treatment	Panicle length* (cm)	Plant fresh weight* (g)	Grains per panicle*	Test weight* (g)	Yield* (q/ha)
T ₁	20.40 ± 0.36 ^d	186.73 ± 3.02 ^{cd}	58.00 ± 3.25 ^{cd}	18.10 ± 0.26 ^d	38.76 ± 0.32 ^d
T ₂	20.30 ± 0.40 ^d	182.73 ± 1.33 ^{de}	58.53 ± 3.20 ^{cd}	18.03 ± 0.30 ^d	38.13 ± 0.68 ^d
T ₃	25.43 ± 0.61 ^b	194.36 ± 4.53 ^{ab}	67.66 ± 4.12 ^b	22.76 ± 1.10 ^b	42.36 ± 0.23 ^b
T ₄	18.80 ± 0.45 ^e	180.00 ± 2.00 ^{ef}	55.70 ± 1.04 ^{de}	16.80 ± 0.30 ^e	34.76 ± 0.51 ^e
T ₅	18.70 ± 0.60 ^e	175.20 ± 2.00 ^{fg}	55.26 ± 2.70 ^{de}	16.73 ± 0.47 ^e	34.10 ± 0.40 ^e
T ₆	21.90 ± 0.72 ^c	191.06 ± 2.20 ^{bc}	61.70 ± 3.04 ^c	19.30 ± 1.01 ^c	40.90 ± 0.36 ^c
T ₇	17.30 ± 0.45 ^f	173.16 ± 5.52 ^g	52.26 ± 2.75 ^e	14.16 ± 0.97 ^f	31.43 ± 1.20 ^f
T ₈	17.23 ± 0.30 ^f	172.20 ± 2.25 ^g	51.96 ± 3.70 ^e	13.93 ± 0.75 ^f	30.53 ± 0.86 ^f
T ₉	17.36 ± 0.51 ^f	173.53 ± 4.53 ^g	52.70 ± 2.08 ^e	14.56 ± 0.65 ^f	31.13 ± 0.50 ^f
T ₁₀	30.06 ± 2.00 ^a	197.10 ± 2.95 ^a	74.62 ± 3.16 ^a	26.66 ± 0.51 ^a	44.00 ± 0.45 ^a
T ₁₁	11.23 ± 0.41 ^g	164.13 ± 4.10 ^h	41.80 ± 5.14 ^f	10.13 ± 0.35 ^g	22.16 ± 0.66 ^g

Different alphabetical letters on the superscript within the columns indicate significant differences among the treatments according to Duncan's multiple range test at $p \leq 0.05$.

T₁ to T₁₁ in the table correspond to the treatment details mentioned in Table 1.

* Data are the average of three replications, $\pm$ represents standard deviations of the mean.

Table 5 Effect of the microbial consortium on superoxide dismutase (U/ml) and hydrogen peroxide (nMol H₂O₂/g FW) of rice.

Treatment	Superoxide dismutase* (U/ml)	Hydrogen peroxide* (nMol H ₂ O ₂ /g FW)
T ₁	0.556 ± 0.04 ^c	0.544 ± 0.02 ^c
T ₂	0.545 ± 0.01 ^c	0.538 ± 0.012 ^c
T ₃	0.723 ± 0.01 ^a	0.705 ± 0.01 ^a
T ₄	0.456 ± 0.02 ^d	0.440 ± 0.04 ^d
T ₅	0.450 ± 0.003 ^d	0.434 ± 0.02 ^d
T ₆	0.631 ± 0.02 ^b	0.621 ± 0.02 ^b
T ₇	0.224 ± 0.003 ^e	0.236 ± 0.004 ^f
T ₈	0.221 ± 0.002 ^e	0.228 ± 0.007 ^f
T ₉	0.226 ± 0.005 ^e	0.239 ± 0.002 ^{ef}
T ₁₀	0.180 ± 0.04 ^f	0.195 ± 0.01 ^f
T ₁₁	0.113 ± 0.006 ^g	0.130 ± 0.006 ^g

Different alphabetical letters on the superscript within the columns indicate significant differences among the treatments according to Duncan's multiple range test at $p \leq 0.05$.

T₁ to T₁₁ in the table correspond to the treatment details mentioned in Table 1.

* Data are the average of three replications, $\pm$ represents standard deviations of the mean.

correlation coefficients (R) of -0.66 and -0.67 , respectively (Supplementary Figure 2).

Discussion

Basmati is considered 'the pearl of scent', because of its texture, aroma, sweetness and long shelf life, having a distinct heritage and special geographical indication (GI). There has been increasing demand of basmati rice in local, national and international markets⁶. However, due to several biotic stresses, such as the bacterial leaf blight of rice caused by *X. oryzae* pv. *oryzae*, its production and productivity has been declining over time. Management of BLB with agro-chemicals has become inefficient and hazardous due to several consumer, environmental and sustainability-related issues. Resistance of bacteria against the antibiotics is also a concern in using such chemicals²⁴. Fifty eight out of 107 isolates of *Xanthomonas smithii* subsp. *citri* have shown resistance against streptomycin, due to chromosomal mutations. Similarly, *Clavibacter michiganensis* has exhibited resistance against streptomycin because of the production of streptomycin-inactivating enzymes²⁶. These negative effects of chemicals have led to the exploration of several other management strategies, such as biological management using various plant growth-promoting agents such as *T. harzianum*, *T. asperellum* and *P. fluorescens*²⁹. Our research on exploring the effectiveness of the microbial consortium containing *T. harzianum* + *T. asperellum* + *P. fluorescens*, using biocontrol agents alone, along with chemicals, against bacterial leaf blight of rice have shown that, foliar application of streptocycline + copper oxychloride resulted in the maximum reduction of percent incidence, percent intensity and lesion length of bacterial leaf blight by 55.01, 59.39 and 52.19 percent, respectively, and exhibited maximum panicle length, plant fresh weight, number of grains per panicle, test weight and yield by 62.64, 16.73, 43.99, 62.00 and 49.62 percent, respectively. Cupric ions (Cu^{++}), the active ingredient in copper oxychloride, caused denaturation of structural and enzymatic proteins along with alterations in membrane semipermeability, resulting in the death of phytopathogens¹⁴. Additionally, streptocycline produces several antibiotics and enhances host defense by generating inhibitors such as H_2O_2 , siderophores and organic acids. The observed reduction in BLB incidence, intensity, and lesion length following the consortium application can be attributed to several synergistic mechanisms. First, both *Trichoderma* spp. and *P. fluorescens* are well-documented producers of antimicrobial compounds, such as antibiotics, siderophores, and lytic enzymes, which directly antagonize *X. oryzae* pv. *oryzae* (*Xoo*)³¹. *Pseudomonas* strains have been shown to produce siderophores and secondary metabolites, such as phenazines and rhamnolipids, which inhibit *Xoo* growth and disrupt pathogen colonization⁴⁹. Similarly, *Trichoderma* spp. secretes chitinases and glucanases that degrade pathogen cell walls, further limiting disease progression³⁰.

Due to several health hazards and environmental pollution, the current investigations to evaluate the effectiveness of biocontrol agents, either alone or in combination (*T. harzianum* + *T. asperellum* + *P. fluorescens*) were used to inhibit *X. oryzae* pv. *oryzae* causing bacterial leaf blight in

basmati rice. The present study demonstrates that the application of a microbial consortium significantly suppressed bacterial leaf blight (BLB) and enhanced key agronomic parameters in Basmati rice under field conditions. These results are in agreement with recent literature, which highlights the multifaceted mechanisms through which beneficial microorganisms mediate plant disease suppression and growth promotion⁴⁰.

This study revealed that seed treatment using the microbial consortium (*T. harzianum* + *T. asperellum* + *P. fluorescens*) reduced disease incidence by 48.56 percent, intensity by 52.42 percent and lesion length by 45.41 percent, and significantly enhanced panicle length, plant fresh weight, number of grains per panicle, test weight and yield by 55.83, 15.55, 38.23, 55.49 and 47.68 percent, respectively. The microbial consortium expressed multiple mechanisms, such as competition, hyperparasitism, secretion of volatile compounds, plant growth-promoting activity, and induction of defence mechanisms that led to increases in the growth of rice plants¹⁹. Maximum increase of fresh and dry root weight by 82.47 and 94.87 percent at 30 DAT, 64.45 and 66.07 percent at 60 DAT and 55.84 and 54.41 percent at 90 DAT were recorded with the seed treatment involving the microbial consortium. Beneficial microbes colonize the root of rice, helping in the proliferation of the roots and shoots of the host, which stimulates plant growth⁷. Applying *T. harzianum* to the rhizosphere is known to expand the root surface area, allowing the roots to explore a greater soil volume and absorb more nutrients. This enhanced nutrient uptake leads to substantial improvements in the growth and yield of rice⁵⁰. Applying *P. fluorescens* (PDY7) has shown better results with a maximum number of grains per panicle of 18.28g as compared to the control (9.11g)⁴⁷. *Bacillus cereus* has shown maximum increase in fresh root weight of rice by 31.4 percent and the application of the microbial consortium (*T. harzianum* + *P. fluorescens* + herbal kunapajala) resulted in the maximum dry root weight of rice³⁶. These enhancements are likely mediated by the plant growth-promoting properties of the constituent microorganisms. Both *Trichoderma* and *Pseudomonas* spp. are known to solubilize inorganic phosphate, produce phytohormones, such as indole acetic acid, and facilitate nutrient uptake, leading to improved root and shoot development³³. The increased root biomass observed in this study is consistent with reports that microbial inoculants enhance root architecture, enabling better water and nutrient acquisition and contributing to overall plant vigor⁴⁰.

In addition to direct antagonism, the microbial consortium appears to induce systemic resistance in rice plants. This is evidenced by the significant increase in superoxide dismutase (SOD) activity and hydrogen peroxide (H_2O_2) concentration observed in consortium-treated plants²³. Our study revealed that seed treatment using the microbial consortium enhanced the amount of superoxide dismutase and hydrogen peroxide by 81.58 and 84.29 percent, respectively. SOD is the reason for detoxification of intercellular O_2^- and secretion of H_2O_2 , removing the dangerous reactive oxygen species (ROS) which results in reduced fitness and fungus virulence^{5,17}. H_2O_2 is sensitive to pathogens but not to the host because H_2O_2 strengthens the cell wall by aiding in the process of lignification, contributing to the cross-linking of cell wall polymers and proline or hydroxyproline-rich

proteins during pathogen invasion, suppressing the growth and development of the invading pathogen²⁰. Previous studies have shown that inoculation with plant growth-promoting rhizobacteria (PGPR) such as *Pseudomonas* spp. can enhance the activity of defense-related enzymes, including peroxidase, catalase, and phenylalanine ammonia-lyase, thereby conferring increased resistance to BLB²⁸. The use of a microbial consortium offers advantages over single-strain inoculants. The combination of multiple beneficial microbes ensures functional redundancy and complementary modes of action, which can buffer against environmental variability and enhance overall effectiveness. This is supported by field studies demonstrating that co-inoculation with multiple antagonistic bacteria or fungi results in greater disease suppression and yield improvement compared to individual strains¹¹.

In the correlation analysis, disease intensity and lesion length exhibited a significantly positive correlation with the incidence of bacterial leaf blight (BLB) of rice. Our findings are in line with earlier studies showing similar results³⁹. Conversely, panicle length, plant fresh weight, fresh root weight at 30, 60 and 90 days after transplanting (DAT), dry root weight at 30, 60, and 90 DAT, number of grains per panicle, test weight, and yield, all showed a negative correlation with BLB incidence, as reported by earlier studies¹³. The application of the microbial consortium increased the uptake of nutrients, the production of siderophores and phytohormones, such as auxins and IAA, resulting in disease mitigation and plant growth promotion³⁷. The correlation analysis of hydrogen peroxide (H₂O₂) concentration and superoxide dismutase activity (SODs) with the incidence of bacterial leaf blight revealed a significantly negative correlation. Biocontrol agents are known to trigger induced systemic resistance by increasing the activity of several enzymes SOD and H₂O₂, PPO and CAT¹.

Conclusion

Seed treatment using the microbial consortium (*T. harzianum* + *T. asperellum* + *P. fluorescens*) proved its effectiveness by reducing percent disease incidence, percent intensity and lesion length of BLB, simultaneously enhancing shoot and root weights, yield and biochemical parameters of rice. These results encourage the use of the microbial consortium for increasing the production of basmati rice by organic means. There is an urgent need for further research to explore the effectiveness of the microbial consortium against several other Basmati rice diseases in large-scale field experiments.

ORCID ID

Fayaz Ahmad Mohiddin: 0000-0001-9720-4429

Funding

The researchers would like to thank the Deanship of Graduate Studies and Scientific Research at Qassim University, for financial support (QU-APC-2024-9/1).

Conflict of interest

The authors declare no conflict of interest.

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version available at <https://doi.org/10.1016/j.ram.2025.08.003>.

References

1. Abdelkhalek A, Aseel DG, Király L, Künstler A, Moawad H, Al-Askar AA. Induction of systemic resistance to Tobacco mosaic virus in tomato through foliar application of *Bacillus amyloliquefaciens* Strain TBorg1 culture filtrate. *Viruses*. 2022;14:1830.
2. Abid J, Ahmed S, Xia T, Wang M. Rice as a nutritional grain: examining its role in healthy products and disease prevention. *Food Rev Int*. 2024;40:3055–78.
3. Adhikari TB, Cruz C, Zhang Q, Nelson R, Skinner D, Mew T, Leach J. Genetic diversity of *Xanthomonas oryzae* pv. *oryzae* in Asia. *AEM*. 1995;61:966–71.
4. Ahmad F, Zhu D, Sun J. Environmental fate of tetracycline antibiotics: degradation pathway mechanisms, challenges, and perspectives. *Environ Sci Eur*. 2021;33:64.
5. Alfei S, Schito GC, Schito AM, Zuccari G. Reactive oxygen species (ROS)-mediated antibacterial oxidative therapies: available methods to generate ROS and a novel option proposal. *Int J Mol Sci*. 2024;25:7182.
6. Amin T, Gupta V, Sharma A, Rai PK, Razdan VK, Sharma SK, Singh SK, Lone JA, Yaqoob M, Singh B. Distribution of *Xanthomonas oryzae* pv. *oryzae* pathotypes in basmati-rice-growing areas of Jammu and Kashmir, India. *Agronomy*. 2023;13:713.
7. Arguelles-Arias A, Ongena M, Halimi B, Lara Y, Brans A, Joris B, Fickers P. *Bacillus amyloliquefaciens* GA1 as a source of potent antibiotics and other secondary metabolites for biocontrol of plant pathogens. *Microb Cell Fact*. 2009;8:1–12.
8. Arif I, Batool M, Schenk PM. Plant microbiome engineering: expected benefits for improved crop growth and resilience. *Trends Biotechnol*. 2020;38:1385–96.
9. Babbal, Adivitiya, Khata YP. Microbes as biocontrol agent. In: *Probiotics and plant health*; 2017. p. 507–52.
10. Baranova N, Sorokin L. Modern educational technology: processing and visualization of the data using the mathematical package originpro 8.5.2 (the concept OF PSYCHO-cognitive barriers). *J Inf Syst Educ*. 2017;14:324–33.
11. Barbosa JZ, Hungria M, Prior SA, Moura MC, Poggere G, Motta ACV. Improving yield and health of legume crops via co-inoculation with rhizobia and *Trichoderma*: a global meta-analysis. *Appl Soil Ecol*. 2022;176, 104493.
12. El-Maraghy SS, Tohamy TA, Hussein KA. Role of plant-growth promoting fungi (PGPF) in defensive genes expression of *Triticum aestivum* against wilt disease. *Rhizosphere*. 2020;15, 100223.
13. Gangwar R, Thorat S, Parmar M, Patel S. Screening of rice genotypes against bacterial leaf blight. *Souvenir Cum Abstracts/Proceeding Book*. Uttarakhand, India: Agricultural & Environmental Technology Development Society; 2021.
14. Grzanka M, Sobiech Ł, Filipczak A, Danielewicz J, Jajor E, Horoszkiewicz J, Korbas M. The efficacy effectiveness of plant pathogens control by complexed forms of copper. *Agriculture*. 2024;14:139.
15. Gupta G, Chauhan PS, Jha PN, Verma RK, Singh S, Yadav VK, Sahoo DK, Patel A. Secretory molecules from secretion systems fine-tune the host-beneficial bacteria (PGPRs) interaction. *Front Microbiol*. 2024;15, 1355750.

16. Haque MM, Masud MM, Bashir S, Hossain MI, Alam MZ, Islam MR. Avirulence gene based RFLP and rep-PCR distinguish the genetic variation of *Xanthomonas oryzae* pv. *oryzae* pathotypes in Bangladesh. *Plant Sci Today*. 2022;9:29–40.
17. Iqbal N, Czékus Z, Ördög A, Poór P. Fusaric acid-evoked oxidative stress affects plant defence system by inducing biochemical changes at subcellular level. *Plant Cell Rep*. 2024;43:2.
18. Jabeen R, Rahman SU, Rais A. Evaluating BLB resistance/aggressiveness in rice through best inoculum concentration, inoculation and application methods. *Pak J Bot*. 2011;43:635–8.
19. Kabdwal BC, Sharma R, Kumar A, Kumar S, Singh K, Srivastava R. Efficacy of different combinations of microbial biocontrol agents against sheath blight of rice caused by *Rhizoctonia solani*, Egypt. *J Biol Pest Control*. 2023;33:29.
20. Kaur S, Samota MK, Choudhary M, Choudhary M, Pandey AK, Sharma A, Thakur J. How do plants defend themselves against pathogens – biochemical mechanisms and genetic interventions. *Physiol Mol Biol Plants*. 2022;28:485–504.
21. Khaskhali S, Xiao X, Solangi F, Hussain S, Solangi KA, Chen Y. Exploring the functional and regulatory mechanisms of the RBOHD gene in rice against bacterial blight disease. *J Plant Interact*. 2025;20, 2456572.
22. Kumar D, Arya SK, Srivastava D, Shamim M, Desai L, Tyagi M. Impact of major rice bacterial diseases on agriculture and food security. In: *Bacterial diseases of rice and their management*. Apple Academic Press; 2023. p. 1–28.
23. Kumar S, Chandra R, Behera L, Keswani C, Sansinenea E. Dual *Trichoderma* consortium mediated elevation of systemic defense response against early blight in potato. *Eur J Plant Pathol*. 2022;162:681–96.
24. Larsson D, Flach CF. Antibiotic resistance in the environment. *Nat Rev Microbiol*. 2022;20:257–69.
25. Louca S, Polz MF, Mazel F, Albright MB, Huber JA, O'Connor MI, Ackermann M, Hahn AS, Srivastava DS, Crowe SA. Function and functional redundancy in microbial systems. *Nat Ecol Evol*. 2018;2:936–43.
26. Lyu Q, Bai K, Kan Y, Jiang N, Thapa SP, Coaker G, Li J, Luo L. Variation in streptomycin resistance mechanisms in *Clavibacter michiganensis*. *Phytopathology*. 2019;109:1849–58.
27. Malabadi RB, Kolkar K, Chalannavar R. White, and brown rice – nutritional value and health benefits: arsenic toxicity in rice plants. *Int J Innov Sci Res Rev*. 2022;4:3065–82.
28. Meena M, Swapnil P, Divyanshu K, Kumar S, Harish, Tripathi YN, Zehra A, Marwal A, Upadhyay RS. PGPR-mediated induction of systemic resistance and physiochemical alterations in plants against the pathogens: current perspectives. *J Basic Microbiol*. 2020;60:828–61.
29. Ngalimat MS, Mohd Hata E, Zulperi D, Ismail SI, Ismail MR, Mohd Zainudin NA, Saidi NB, Yusof MT. A laudable strategy to manage bacterial panicle blight disease of rice using biocontrol agents. *J Basic Microbiol*. 2023;63:1180–95.
30. Parveen S, Mohiddin FA, Bhat MA, Baba ZA, Jeelani F, Bhat MA, Nabi SU, Hamid B, Bandey S, Rasool F. Characterization of endophytic microorganisms of rice (*Oryza sativa* L.) potentials for blast disease biocontrol and plant growth promoting agents. *Phyton*. 2023;92, 0031-9457.
31. Pérez-Pizá MC, Sautua FJ, Szparaga A, Bohata A, Kocira S, Carmona MA. New tools for the management of fungal pathogens in extensive cropping systems for friendly environments. *Crit Rev Plant Sci*. 2024;43:63–93.
32. Prodhan ZH, Samonte SOP, Sanchez DL, Talukder SK. Profiling and improvement of grain quality traits for consumer preferable basmati rice in the United States. *Plants*. 2024;13:2326.
33. Rawat P, Das S, Shankhdhar D, Shankhdhar S. Phosphate-solubilizing microorganisms: mechanism and their role in phosphate solubilization and uptake. *J Soil Sci Plant Nutr*. 2021;21:49–68.
34. Resti Z, Liswarni Y, Martinius M. Endophytic bacterial consortia as biological control of bacterial leaf blight and plant growth promoter of rice (*Oryza sativa* L.). *J Appl Agric Sci Technol*. 2020;4:134–45.
35. Saad A, Habibuddin H. Pathotypes and virulence of *Xanthomonas oryzae* causing bacterial blight disease of rice in Peninsular Malaysia. *J Trop Agric Food Sci*. 2010;38:257–66.
36. Sagar S, Singh A, Bala J, Chauhan R, Kumar R, Bhatia RK, Walia A. Insights into cow dung-based bioformulations for sustainable plant health and disease management in organic and natural farming system: a review. *J Soil Sci Plant Nutr*. 2024;24:30–53.
37. Seenivasagan R, Babalola OO. Utilization of microbial consortia as biofertilizers and biopesticides for the production of feasible agricultural product. *Biology*. 2021;10:1111.
38. SES: standard evaluation system for rice. Philippines: International Rice Research Institute; 2013.
39. Shah SMA, Ahsan R, Liu L, Li Y, Wang Q, Wang Y, Yan J, Khojasteh M, Xu X, Xu Z. TALome and phenotypic analysis of Pakistani *Xanthomonas oryzae* pv. *oryzae* population revealed novel virulent TALEs contributing to bacterial blight of rice. *Phytopathol Res*. 2025;7:5.
40. Sharma A, Gupta V, Jamwal G, Gupta SK, Pandit V, Amin Z, Mohiddin FA, Hamed KE, Ashraf S. Harnessing microbial consortia for sustainable management of corm rot of saffron. *J Basic Microbiol*. 2025;65:e70009.
41. Shekhar S, Sinha D, Kumari A. An overview of bacterial leaf blight disease of rice and different strategies for its management. *Int J Curr Microbiol Appl Sci*. 2020;9:2250–65.
42. Singh DP, Singh V, Gupta VK, Shukla R, Prabha R, Sarma BK, Patel JS. Microbial inoculation in rice regulates antioxidative reactions and defense related genes to mitigate drought stress. *Sci Rep*. 2020;10:4818.
43. Suryadi Y, Samudra I, Priyatno TP, Susilowati DN, Lestari P, Kadir TS. Determination of pathotypes from Indonesian *Xanthomonas oryzae* pv. *oryzae* population causing bacterial leaf blight and their reactions on differential rice. *Makara J Sci*. 2016;20:3.
44. Swings J, Van den Mooter M, Vauterin L, Hoste B, Gillis M, Mew T, Kersters K. Reclassification of the causal agents of bacterial blight (*Xanthomonas campestris* pv. *oryzae*) and bacterial leaf streak (*Xanthomonas campestris* pv. *oryzicola*) of rice as pathogens of *Xanthomonas oryzae* (ex Ishiyama 1922) sp. nov., nom. rev. *Int J Syst Evol Microbiol*. 1990;40:309–11.
45. Tekete C, Cunnac S, Doucouré H, Dembele M, Keita I, Sarra S, Dagno K, Koita O, Verdier V. Characterization of new races of *Xanthomonas oryzae* pv. *oryzae* in Mali informs resistance gene deployment. *Phytopathology*. 2020;110:267–77.
46. Trivedi S, Srivastava M, Ratan V, Mishra A, Dixit S, Pandey S. Evaluation of microbial consortia on systemic resistance against chickpea wilt. *Bangladesh J Bot*. 2020;49:653–61.
47. Velusamy P, Immanuel JE, Gnanamanickam SS. Rhizosphere bacteria for biocontrol of bacterial blight and growth promotion of rice. *Rice Sci*. 2013;20:356–62.
48. Westfall PH, Arias AL, Fulton LV. Teaching principal components using correlations. *Multivar Behav Res*. 2017;52:648–60.
49. Yang R, Li S, Li Y, Yan Y, Fang Y, Zou L, Chen G. Bactericidal effect of *Pseudomonas oryziphila* sp. nov., a novel *Pseudomonas* species against *Xanthomonas oryzae* reduces disease severity of bacterial leaf streak of rice. *Front Microbiol*. 2021;12, 759536.
50. Yedidia I, Srivastava AK, Kapulnik Y, Chet I. Effect of *Trichoderma harzianum* on microelement concentrations and increased growth of cucumber plants. *Plant Soil*. 2001;235:235–42.